

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

verse results of the East Indian experiments with C. panamensis may not apply to the whole genus. Moreover, during the present study of the subject many reasons have been found for believing that the conditions under which Castilla has been tested in the East Indies are not really favorable to the production of rubber; the current idea that a continuously humid climate is required is erroneous. In short, it appears that we are still at the beginning of a scientific comprehension of the factors which determine the practicability and profitability of rubber culture. It has been ascertained that rubber can be produced agriculturally, but where, how and what to plant, and how, how much and how long we shall harvest, are questions largely answered, as yet, by speculation rather than by experiment. O. F. Cook.

U. S. DEPARTMENT OF AGRICULTURE.

THE NAME OF THE BREADFRUIT.

The genus Artocarpus was first described in 1776 by G. and G. J. R. Forster in the 'Characteres Generum Plantarum,' a work written as a result of their botanical studies made during Captain Cook's second voyage into the Pacific and round the world between 1772 and 1775. The combination Artocarpus communis was given in this work for the breadfruit tree, a name which, according to nomenclatorial rules, must replace the generally accepted Artocarpus incisa, which was not published by the younger Linnæus until 1781.*

Forster's genus was, moreover, published as a monotype, and as his plants were from the Society Islands there can be no doubt but that he was dealing with the true breadfruit. He did not publish, it is true, any specific description, leaving all for the genus, but he did make a good binomial combination and had two good plates which are generally considered sufficient to establish a name in good standing.

Thunberg later in the same year (1776) published the names *Radermachia incisa* and *integrifolia* for the bread- and jak-fruits respectively from material collected in the

East Indian Islands. Five years afterwards the younger Linnæus made his new nomenclatorial combinations on this material of Thunberg, adopting Forster's generic name and adding to it Thunberg's specific designations, and taking the credit to himself.

Further complications are met with when it is found that in the subsequent works of the Forsters, when mention is made of the breadfruit, the specific name *incisa* is used. Why they should abandon their own name is rather difficult to understand unless it was a case in which 'the king can do no wrong.'

Dr. A. Richter is fully alive to the injustice done Forster and has published a note* on the history of the name of the breadfruit which adequately states the facts in the case and further calls attention to the unfortunate revival by O. Kuntze of the pre-Linnæan name of *Soccus*, a relic of Rumphius, and of his combining with it Forster's specific name. Yet Rumphius published a specific name for the breadfruit which Kuntze has, for no apparent reason, seen fit to ignore.

A. Engler, acting on this note, has corrected in the 'Nachtrag' to the 'Natürlichen Pflanzenfamilien' the name of the breadfruit as it appears in the text of that work, and states that *Artocarpus communis* is the correct designation.

Henry E. Baum.

U. S. DEPT. AGRICULTURE.

EUCALYPTS IN THE PHILIPPINES.

The eucalypts, of which but comparatively few species are familiarly known outside of their native home, include some one hundred and fifty species or more, nearly all restricted to Australia and Tasmania. Many of the forms may be classed as shrubs, others attain great size, surpassing in height, as has been stated on good authority, the giant Sequoias of California, though not equaling them in diameter or girth. A few species have been found elsewhere, viz., in New Britain, New Guinea and Timor, islands north of the Australian continent, between latitude 10° S. and the equator. It is not unlikely that sooner or later other species, at present unknown, will be detected on some of the multi-

^{* &#}x27;Suppl.' 411. 1781.

^{*} Botanisches Centralblatt 60: 169-170. 1894.